



# YUVEZZI ECP Toolkit FAQ

## PURPOSE

This document provides prepared responses for common questions that you can deliver in the event you receive inbound inquiries from members of the media about YUVEZZI™. Please ensure you refer to the Important Safety Information below and provide fair balance of safety information in your responses. Please note that these messages are not to be distributed and must be used exactly as instructed. Share the patient counseling information, included below, with patients at the time of your interactions.

## FAQ

### What is YUVEZZI?

- YUVEZZI is a prescription eye drop for adults with blurry close-up vision due to age (presbyopia).
- YUVEZZI is the first and only dual-agent (combination) eye drop approved for the treatment of presbyopia in adults.

### How does YUVEZZI work?

- YUVEZZI combines two active ingredients that work together to treat age-related blurry close-up vision (presbyopia).
  - Carbachol helps your pupils get smaller, or constrict, so it's easier to focus up close.
  - Brimonidine helps keep your pupils from getting too large, or dilating.
- Both active ingredients in YUVEZZI have been trusted to treat other common eye conditions for nearly 30 years.

### How do I take YUVEZZI?

- YUVEZZI is taken once daily.
- You put 1 drop of YUVEZZI in each eye, every day. Twist off the top of the plastic vial, squeeze the vial to release 1 drop in each eye and throw away the vial (even if it's not empty). If you wear contact lenses, remove them before using YUVEZZI. Wait at least 10 minutes after using YUVEZZI before you put them back in your eyes. Be sure to not let the tip of the single-use vial touch your eye, eyelid, or any other surface.

### Will YUVEZZI replace my reading glasses?

- YUVEZZI can be used your way: on its own instead of reading glasses or with contacts or glasses you use to see far away.
- Note: If you wear contact lenses, remove them before using YUVEZZI. Wait at least 10 minutes after using YUVEZZI before you put them back in your eyes.

### Can I use YUVEZZI if I wear glasses or contacts for distance vision?

- Yes, YUVEZZI can be used with contacts or glasses that you use to see far away.
- Note: If you wear contact lenses, remove them before using YUVEZZI. Wait at least 10 minutes after using YUVEZZI before you put them back in your eyes.

### How long does it take to work?

- In medical studies, as soon as 30 minutes after taking YUVEZZI, some people could clearly read at least 3 more lines on an eye chart in low light without losing distance vision.

### How long does it last?

- In medical studies, YUVEZZI provided up to 8 continuous hours of improved close-up vision for about half of the people. And YUVEZZI worked just as well over time: after 12 months of use, YUVEZZI once-daily drops consistently improved blurry close-up vision like it did at the start.

### Are there any side effects?

- In a medical study of YUVEZZI in 358 patients, the most common side effects were headache (16%), impaired vision (6%), and eye pain (7%), and/or eye irritation upon use (14%).
  - Most side effects were generally mild, didn't last long, and went away on their own
  - Headache rates were generally low (less than 2 of 10 patients) and mostly stopped after the first week of use
  - People taking YUVEZZI had no treatment-related serious side effects in a 1-year safety study
  - Nearly 9 out of 10 people were able to keep using YUVEZZI without stopping because of side effects for the full year of the study
- These are not all of the possible side effects of YUVEZZI.

### Does it need to be refrigerated?

- No, YUVEZZI does not need to be refrigerated.

### How much does YUVEZZI cost?

- For the most up-to-date pricing information, visit [www.yuvezzi.com](http://www.yuvezzi.com).

---

### Patient Counseling Information.

YUVEZZI is contraindicated in patients with known hypersensitivity to the active ingredients or to any of the ingredients.

#### Night Driving

Patients should exercise caution when night driving, as YUVEZZI may cause temporary dim or dark vision.

#### Blurred Vision

Temporary problems when changing focus between near objects and distant objects may occur.

#### When to Seek Physician Advice

Advise patients to seek immediate medical care with sudden onset of flashes of lights, floaters, or vision loss.

#### Contact Lens Wear

Contact lenses should be removed prior to the instillation of YUVEZZI. Wait 10 minutes after dosing before reinserting contact lenses.

#### Concomitant Topical Ocular Therapy

If more than one topical ophthalmic medication is being used, the medicines must be administered at least 5 minutes apart.

#### Avoiding Contamination of the Product

Patients should avoid touching the tip of the single-use vial to the eye or to any other surface.

For additional safety information, please visit the YUVEZZI [Brief Summary](#).

---

## USE

YUVEZZI (carbachol and brimonidine tartrate ophthalmic solution) 2.75% / 0.1% is a prescription eye drop for adults with blurry close-up vision due to age.

## IMPORTANT SAFETY INFORMATION

Do not use YUVEZZI if you are allergic to any of its ingredients or if you currently have inflammation of the iris (iritis).

Before taking YUVEZZI, tell your doctor if you have depression, low blood pressure, or circulation problems.

YUVEZZI may cause temporary blurry, dim, or dark vision. If you experience this, avoid driving, using machinery, and participating in hazardous activities. Use caution when night driving and in other activities in low light.

Call your doctor right away if you suddenly have flashes of light, floaters, or vision loss.

Do not let the tip of the vial touch your eye, eyelid, or any other surface.

The most common side effects of YUVEZZI are headache, impaired vision, and eye pain and/or eye irritation upon use. These are not all of the possible side effects of YUVEZZI.

Most side effects were generally mild, didn't last long, and went away on their own.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Please see the [Brief Summary](#) for more information.

